

Monroe County Influenza Surveillance October 1, 2013 –March 29, 2014 Rochester - Emerging Infections Program (EIP) Center for Community Health-URMC

(Data summary provided to Monroe County Department of Public Health)


This is a peliminary report of Influenza cases. As of 3/29/2014, there were 1546 laboratory confirmed Influenza cases (1511 Flu A and 35 Flu B) reported and 335 hospitalizations



Laboratory Surveillance: Positive influenza tests are reported on the NY State Health Information Network (HIN) by all diagnostic laboratories. The above figure shows the number and type of positive tests from Monroe County residents, and the percent of all tests performed that were positive for flu.




Emergency Room Surveillance: All 4 Monroe County Hospitals report to the HIN the total number of ED visits each day. Influenza-like illness (ILI) cases are those that have fever or fever and respiratory symptoms as chief complaint.



Long Term Care (LTCF) Surveillance: Beginning October 1, 2013, 32 Monroe County LTCFs are asked to voluntarily report ILI and flu activity each week.

Monroe County Influenza Surveillance October 1, 2013 –March 29, 2014 Rochester - Emerging Infections Program (EIP) Center for Community Health-URMC

(Data summary provided to Monroe County Department of Public Health)



Number of hospitalizations due to community acquired and hospital onset influenza cases as of **March 29, 2014**. Hospitalization data is collected up to 14 days of positive influenza test. Definitions detailed below:

- Community acquired influenza hospitalization includes patients who were hospitalized **14 days or less** *after* a positive influenza test <u>or</u> **3 days or less** *before* a positive influenza test
- Hospital onset influenza cases include patients who have a positive influenza test after 3 days of admission.

Cumulative Number of Admissions and Deaths in Monroe County October 1, 2013 to March 29, 2014

Age Group	# Cases Hospitalized N (%)	Deaths N (%)
0 to 4 y	13 (4%)	0 (%)
5 to 17 y	12 (3%)	0 (%)
18 to 49 y	112 (34%)	5 (36%)
50 to 64 y	111 (33%)	5 (36%)
≥ 65 y	87 (26%)	4 (28%) *
Total	335 (100%)	14 (100%)

* One of the deaths occurred in the ≥ 65 age group was a non-hospitalized patient